


IN THE CLAIMS

Please amend the claims as follows:

Claim 1 (Currently Amended): A compound having the formula (1):



wherein

R^1 is selected from the group consisting of H, NO_2 , CN, OCH_3 , a halogen, an alkyl having up to 4 carbon atoms, and an alkoxy having up to 4 carbon atoms;


R^2 is selected from the group consisting of an aryl group, a substituted aryl group, a heteroaryl group, substituted heteroaryl group, an aroyl group, and a substituted aroyl group;

R^3 is selected from the group consisting of H, NO_2 and a halogen;


R^4 is selected from the group consisting of H, OCH_3 and an alkyl group having up to 4 carbon atoms;

X is selected from the group consisting of oxygen and sulfur; and

Z is selected from the group consisting of a leaving group, an alcoholate group, $-OH$, a N-atom of an amine compound, a deoxyribonucleoside and a ribonucleoside as represented by either of the following formulae (2) or (3):



2



3

wherein

R^5 is selected from the group consisting of a H, an oligonucleotide, a phosphitamidite group and a protecting group functional group useful in oligonucleotide synthesis;

R^6 is selected from the group consisting of H, OH, an alkoxyl having up to 4 carbon atoms, an alkenoxyl group having up to 4 carbon atoms, a substituted alkenoxyl group having up to 4 carbon atoms, or R^6 is WR^8 wherein W is selected from oxygen and sulfur and R^8 is a protective group useful in oligonucleotide synthesis;

B is a base selected from the group consisting of adenine, cytosine, guanine, thymine, and uracil and chemical modifications thereof, and when B is any one of adenosine, cytosine and guanine the amino functions on the heterocycle may bear a protective group useful in oligonucleotide synthesis; ~~or~~

~~Z is selected from the group consisting of a chemically modified deoxyribonucleoside, a chemically modified ribonucleoside, and an analog thereof.~~

Claim 2 (Canceled).

Claim 3 (Previously Presented): The compound of claim 1, wherein R^1 is H and R^2 is phenyl or substituted phenyl.

Claim 4 (Previously Presented): The compound of claim 1, wherein R^1 is H and R^2 is benzoyl or substituted benzoyl.

Claim 5 (Previously Presented): The compound of claim 1 wherein W is O and R^8 is selected from the group consisting of an alkyl, alkenyl, acetal and silylether protective group.

Claim 6 (Previously Presented): The compound of claim 1, wherein W is S and R^8 is an alkyl protective group.

Claim 7 (Previously Presented): The compound of claim 1, wherein R^6 is selected from the group consisting of an O-methyl, O-ethyl, O-allyl, O-tetrahydropyranyl- O-methoxytetrahydropyranyl and an O-t-butyl dimethylsilyl.

Claim 8 (Previously Presented): The compound of claim 1, wherein B is selected from the group consisting of adenine, cytosine and guanine and wherein R^8 is selected from the group consisting of phenoxyacetyl, 4-tert-butyl-phenoxyacetyl, 4-isopropyl-phenoxyacetyl and dimethylformamidino.

Claim 9 (Previously Presented): The compound of claim 1, wherein B is adenine and is selected from the group consisting of benzoyl and p-nitrophenyloxycarbonyl (p-NPEOC).

Claim 10 (Previously Presented): The compound of claim 1, wherein B is guanine and wherein R^8 is selected from the group consisting of isobutyroyl and p-nitrophenylethyloxycarbonyl (p-NPEOC).

Claim 11 (Previously Presented): The compound of claim 1, wherein B is cytosine and wherein R⁸ is selected from the group consisting of benzoyl, isobutyryl and p-nitrophenylethoxy carbonyl (p-NPEOC).

Claim 12 (Currently Amended): The compound of claim 1, wherein R⁵ is a ~~phosphitamide~~ phosphitamidite group.

Claim 13 (Previously Presented): The compound of claim 1, wherein R⁵ is an OH-protective group.

Claim 14 (Previously Presented): The compound of claim 13, wherein R⁵ is a dimethoxytrityl- or a monomethoxytrityl- group.

Claim 15 (Original): The compound of claim 13, wherein R⁵ is a silyl-group.

Claim 16 (Previously Presented): The compound of claim 1, wherein Z is a leaving group.

Claim 17 (Previously Presented): The compound of claim 16, wherein the leaving group is selected from the group consisting of chloride, imidazolyl and nitrophenoxyl.

Claims 18-23 (Canceled)

Claim 24 (Withdrawn-Currently Amended): A method for the light-controlled synthesis of oligonucleotides, wherein said method is comprised of the following steps:

a) attaching, as a first building block, a nucleoside or nucleotide of claim 1 comprising the photolabile protective group at its primary hydroxyl group, to a support via its 3' secondary hydroxyl group;

b) irradiating the support-bound nucleoside or nucleotide resulting from step a), such that the protective group at the primary hydroxyl group is removed, thereby deprotecting the primary hydroxyl group;

c) reacting the support-bound nucleotide resulting from step b) in the presence of an activator with a second nucleotide selected from claim 12 comprising a protective group at its primary hydroxyl group and phosphoramidite functional group at its 3' secondary hydroxyl group, to form an internucleosidic phosphorous linkage;

d) optionally capping unreacted primary hydroxyl groups with an inert alcohol protecting group;

e) oxidizing the internucleosidic phosphorous linkage to the naturally occurring pentavalent state;

f) iterating steps b) to d) while successively applying the phosphoramidite building blocks in a predetermined order until the desired oligonucleotide strand is completed; and

g) removing of all nucleobase and phosphate protective groups.


Claim 25 (Withdrawn-Currently Amended): A method for the light-controlled synthesis of oligonucleotides, wherein said method is comprised of the following steps:

a) attaching, [[a]] as a first building block, a nucleoside or nucleotide of claim 1 comprising the photolabile protective group at its 3' secondary hydroxyl group, to a support via its primary hydroxyl group;

- b) irradiating the support-bound nucleotide resulting from step a), such that the protective group at the secondary hydroxyl group is removed, thereby deprotecting the 3' secondary hydroxyl group;
- c) reacting the support-bound nucleotide resulting from step b) in the presence of an activator with a second nucleotide selected from claim 12 comprising a protective group at its 3' secondary hydroxyl group and a phosphoramidite functional group at its primary hydroxyl group, to form an internucleosidic phosphorous linkage;
- d) optionally capping unreacted secondary hydroxyl groups with an inert alcohol protecting group;
- e) oxidizing the internucleosidic phosphorous linkage to the naturally occurring pentavalent state;
- f) iterating steps b) to d) while successively applying the phosphoramidite building blocks in a predetermined order until the desired oligonucleotide strand is completed; and
- g) removing of all nucleobase and phosphate protective groups.

Claims 26-29 (Canceled).

Claim 30 (Currently Amended): A compound having the formula (1):



wherein

R^1 is $COOY$, wherein Y is selected from the group consisting of an alkyl group of up to 10 carbon atoms,



R^2 is selected from the group consisting of H, NO_2 , CN, OCH_3 , a halogen, an alkyl having up to 4 carbon atoms, an alkoxy having up to 4 carbon atoms;

R^3 is selected from the group consisting of H, NO_2 and halogen;

R^4 is selected from the group consisting of OCH_3 , an alkyl group having up to 4 carbon atoms and an optionally substituted alkyl group having up to 4 carbon atoms;

X is selected from the group consisting of oxygen or sulfur; and

Z is selected from the group consisting of a leaving group, an alcoholate, $-OH$, a N-atom of an amine compound, a deoxyribonucleoside and a ribonucleoside as represented by either of the following formulae (2) or (3):



wherein

R^5 is selected from the group consisting of a H, an oligonucleotide, a phosphitamidite group and a protecting group functional group useful in oligonucleotide synthesis;

R^6 is selected from the group consisting of H, OH, an alkoxy having up to 4 carbon atoms, an alknoxy having up to 4 carbon atoms, or a substituted alkenoxy having up to 4 carbon atoms, or R^6 is WR^8 wherein W is selected from oxygen and sulfur and R^8 is selected from a protective group useful in oligonucleotide synthesis;

B is base selected from the group consisting of adenine, cytosine, guanine, thymine, and uracil, and chemical modifications thereof and in the case of adenosine, cytosine and

guanine the amino functions on the heterocycle may bear a protective group useful in oligonucleotide synthesis; or

~~Z is selected from the group consisting of a chemically modified deoxyribonucleoside, a chemically modified ribonucleoside, and an analog thereof.~~

Claim 31 (Previously Presented): The compound of claim 30, wherein Y is an alkyl group selected from the group consisting of methyl and tertiary-butyl, and R² is H.

Claim 32 (Previously Presented): The compound of claim 30 wherein W is O and R⁸ is selected from the group consisting of an alkyl, alkenyl, acetal and silyl ether protective group.

Claim 33 (Previously Presented): The compound of claim 30 wherein W is S and R⁸ is selected from the group consisting of an alkyl protective group.

Claim 34 (Previously Presented): The compound of claim 30, wherein R⁶ is selected from the group consisting of an O-methyl, O-ethyl, O-allyl, O-tetrahydropyranyl- O-methoxytetrahydropyranyl and an O-t-butyl dimethylsilyl.

Claim 35 (Previously Presented): The compound of claim 30, wherein B is selected from the group consisting of adenine, cytosine and guanine and said protective group is selected from the group consisting of phenoxyacetyl, 4-tert-butyl-phenoxyacetyl, 4-isopropyl-phenoxyacetyl and dimethylformamidino.

Claim 36 (Previously Presented): The compound of claim 30, wherein B is adenine and the protective group is selected from the group consisting of benzoyl and p-nitrophenyloxycarbonyl (p-NPEOC).

Claim 37 (Previously Presented): The compound of claim 30, wherein B is guanine and the protective group is selected from the group consisting of isobutyryl and p-nitrophenylethyloxycarbonyl (p-NPEOC).

Claim 38 (Previously Presented): The compound of claim 30, wherein B is cytosine and the protective group is selected from the group consisting of benzoyl, isobutyryl and p-nitrophenylethyloxycarbonyl (p-NPEOC).

Claim 39 (Currently Amended): The compound of claim 30, wherein R⁵ is a ~~phosphitamide~~ phosphitamidite group.

Claim 40 (Previously Presented): The compound of claim 30, wherein R⁵ is an OH-protective group.

Claim 41 (Previously Presented): The compound of claim 40, wherein R⁵ is selected from a dimethoxytrityl- or a monomethoxytrityl- group.

Claim 42 (Previously Presented): The compound of claim 40, wherein R⁵ is a silyl-group.


Claim 43 (Previously Presented): The compound of claim 30, wherein Z is a leaving group.

Claim 44 (Previously Presented): The compound of claim 43, wherein the leaving group is selected from the group consisting of chloride, imidazolyl and nitrophenoxyl.


Claim 45 (Previously Presented): The compound of claim 31, wherein Z is a leaving group.

Claim 46 (Previously Presented): The compound of claim 45, wherein the leaving group is selected from the group consisting of chloride, imidazolyl and nitrophenoxyl.

Claim 47 (Previously Presented): The compound of claim 1, wherein Z is a dcoxyribonucleoside or a ribonucleoside represented by formulae (2) or (3):



2



3

wherein R⁵ is selected from the group consisting of a H and an oligonucleotide;

R⁶ is selected from the group consisting of H, OH, an alkoxy having up to 4 carbon atoms, an alkenoxy group having up to 4 carbon atoms, a substituted alkenoxy group having up to 4 carbon atoms, or R⁶ is WR⁸ wherein W is selected from oxygen and sulfur and R⁸ is a protective group;

B is selected from the group consisting of adenine, cytosine, guanine, thymine and uracil.

Claim 48 (Previously Presented): The compound of claim 1, wherein Z is selected from the group consisting of a deoxyribonucleoside and a ribonucleoside.

Claim 49 (Previously Presented): The compound of claim 1, wherein Z is selected from the group consisting of an alcoholate group, -OH and an amine.


Claim 50 (Previously Presented): The compound of claim 1, wherein

R² is a phenyl group;


R⁴ is a hydrogen atom or an alkyl group having up to 4 carbon atoms;

X is O;

and Z is a deoxyribonucleoside or a ribonucleoside represented by formulae (2) or (3):



2




3

wherein R⁵ is H;


R⁶ is selected from the group consisting of H, OH, an alkoxyl having up to 4 carbon atoms, and an alkenoxyl group having up to 4 carbon atoms;
and

B is selected from the group consisting of adenine, cytosine, guanine, thymine and uracil.

Claim 51 (Previously Presented): The compound of claim 30, wherein Z is a deoxyribonucleoside or a ribonucleoside represented by formulae (2) or (3):



2



3

wherein R⁵ is selected from the group consisting of a H and an oligonucleotide;

R⁶ is selected from the group consisting of H, OH, an alkoxy having up to 4 carbon atoms, an alkenoxy having up to 4 carbon atoms, a substituted alkenoxy having up to 4 carbon atoms, or R⁶ is WR⁸ wherein W is selected from oxygen and sulfur and R⁸ is a protective group;

B is selected from the group consisting of adenine, cytosine, guanine, thymine and uracil.

Claim 52 (Previously Presented): The compound of claim 30, wherein Z is selected from the group consisting of a deoxyribonucleoside and a ribonucleoside.


Claim 53 (Previously Presented): The compound of claim 30, wherein Z is selected from the group consisting of an alcoholate group, -OH and an amine.

Claim 54 (Previously Presented): The compound of claim 30, wherein R² is a phenyl group;


R^4 is a hydrogen atom or an alkyl group having up to 4 carbon atoms;

X is O;

and Z is a deoxyribonucleoside or a ribonucleoside represented by formulae (2) or (3):



2



3

wherein R^5 is H;

R^6 is selected from the group consisting of H, OH, an alkoxy having up to 4 carbon atoms, and an alkenoxy having up to 4 carbon atoms;
and

B is selected from the group consisting of adenine, cytosine, guanine, thymine and uracil.

Claim 55 (Previously Presented): The compound of claim 1, wherein R^1 and R^3 are selected from the group consisting of H and NO_2 , wherein R^1 and R^3 are not both NO_2 ;

R^2 is selected from the group consisting of a phenyl group and a benzoyl group;

R^4 is selected from the group consisting of a methyl group and an ethyl group; and

X is oxygen.

Claim 56 (Previously Presented): The compound of claim 55, wherein Z is a deoxyribonucleoside;

R^6 = H; and

B is selected from the group consisting of adenine, cytosine, guanidine, thymine and uracil.

Claim 57 (Previously Presented): The compound of claim 1, wherein Z is a deoxyribonucleoside;

$R^6 = H$; and

B is selected from the group consisting of adenine, cytosine, guanidine, thymine and uracil.